

EXPERIMENTAL VACCINE PROTECTION AGAINST HOMOLOGOUS AND HETEROLOGOUS STRAINS OF FELINE IMMUNODEFICIENCY VIRUS

RUNNING TITLE: Experimental Vaccine Protection Against FIV

**Janet K. Yamamoto,^{ah} Tsutomu Hohdatsu,^{ab} Robert A. Olmsted,^{cd} Ruiyu Pu,^a Harry Louie,^a
Howard A. Zochlinski,^e Veronica Acevedo,^a Howard M. Johnson,^f Gregory A. Soulds,^g and
Murray B Gardner^e**

^a Department of Medicine, MS-1A, School of Veterinary Medicine, University of California, Davis, California 95616.

^b Department of Veterinary Infectious Diseases, School of Veterinary Medicine and Animal Sciences, Kitasato University, Aomori-ken, Japan 034.

^c Division of Molecular Virology and Immunology, Department of Microbiology, Georgetown University, Rockville, Maryland 20852.

^d Present address: Upjohn Laboratories, Cancer & Infectious Disease Research, Kalamazoo, Michigan, 49001.

^e Department of Medical Pathology, School of Medicine, University of California, Davis, California 95616.

^f Department of Microbiology and Cell Sciences, University of Florida, Gainesville, Florida 32611.

^g Synbiotics Corporation, San Diego, California 92127.

^h Corresponding author: Phone number (916) 752-0389



ABSTRACT

The experimental infection of domestic cats with feline immunodeficiency virus (FIV) represents a useful small animal model for testing AIDS vaccine prophylaxis. We previously reported that inactivated whole infected cell or cell-free virus vaccines protected about 75% (11 of 15) immunized cats against intraperitoneal (IP) infection with a low dose of FIV, Petaluma strain. We now report that, using similar but slightly modified FIV vaccine protocols, over 90% of immunized cats were protected against IP infection with the same dose of either homologous FIV (Petaluma strain) (28 of 30 cats) or heterologous FIV (Dixon strain) (27 of 28 cats). These FIV strains differ by 11% in outer envelope amino acid sequences. All control cats (15 of 15) were readily infected with either strain of FIV. Vaccine induced immunity appeared sterilizing and correlated with strong antiviral antibody responses, although not necessarily neutralizing antibody. The adjuvant role of autologous feline T-cellular antigens and duration of immunity in this vaccine model remain to be determined.

Infection of domestic house cats with feline immunodeficiency virus (FIV) is an established cause of fatal immunosuppressive disease under both natural and experimental conditions and is becoming a very useful small animal model for AIDS pathogenesis, antiviral therapy and vaccine testing (13,3). We previously showed that sterilizing immunity against low-dose intraperitoneal (IP) infection with FIV could be achieved in about 75% of cats immunized with inactivated whole FIV vaccines made up of either fixed infected allogenic T-cells or cell-free virus harvested from these cells (20). Protection appeared to correlate in general with vaccine induction of a strong antiviral immune response including neutralizing antibodies. However, the relative titer of neutralizing antibodies did not completely correlate with protection. In contrast to the SIV macaque model (16), immunization of cats with the uninfected T-cells used to grow the virus for vaccine production and challenge, did not protect any cats against challenge infection with FIV (20). Thus, in the FIV model, although the allogeneic T-cell antigens could be shown to contribute an adjuvant effect to the vaccines (20), they did not appear to be the critical correlate of immune protection. In the present study we expanded upon this FIV vaccine model by immunizing a larger number of cats with a slightly modified protocol and by challenging cats protected against the homologous FIV with a heterologous strain of FIV, differing by 11% in outer envelope amino acid sequences.

The IL-2 independent feline T-cell line (FL-4) used for the preparation of the FIV vaccines has been described (18,20). In short, FIV-infected FL-4 cells were fixed in 1.25% paraformaldehyde for 24 hr at 5°C and stored at 5°C (2.5×10^7 cells/ml). Equal volumes (0.5 ml) of the cell suspension and adjuvant, adenylyl muramyl dipeptide (A-MDP) (500 mg/ml) were mixed immediately prior to vaccination. The cell-free virus preparation consisted of filtered (0.45 μ) FL-4 tissue culture fluid pelleted by ultracentrifugation (17000-37000 rpm),

resuspended in phosphate buffered saline (PBS) and treated with 1.25% paraformaldehyde for 24 hr at 5°C. The inactivated virus was resuspended in PBS at a concentration of 500 µg/ml and stored at -70°C. Immediately prior to vaccination, the inactivated virus preparation was thawed at room temperature and mixed with an equal value (0.5 ml) of A-MDP adjuvant. Both fixed-cell and inactivated whole-virus vaccines were shown to be free of infectious virus and viable cells by in vitro virus isolation and proliferation assays and by monitoring vaccinated unchallenged cats for infectious virus using virus isolation and PCR for over 2 years (data not shown).

Forty specific pathogen free (SPF) cats 4-6 months of age were divided into 3 groups: Group A (15 cats) was vaccinated subcutaneously with 2.5×10^7 fixed infected FL-4 cells per dose plus A-MDP adjuvant, Group B (15 cats) received 250 µg of inactivated whole virus per dose plus A-MDP adjuvant and Group C (10 cats) received adjuvant alone. All cats were given the 1st boost 2 weeks after the primary immunization and the 2nd boost 3 weeks later. Because of apparent A-MDP toxicity, i.e., diarrhea, vomiting, following the primary immunization, the dose of A-MDP was reduced to 125 µg and 10 µg for the 1st and 2nd boosts.

Following vaccination but before challenge with homologous virus (day 0, Figs 1-4) all cats were monitored for their humoral immune response and presence or absence of infectious virus in cultured PBMCs. They remained completely negative for infectious virus by both virus isolation and PCR (Table 1A), further evidence that the vaccines were completely inactivated in vitro. By 3 weeks after the 2nd boost with either vaccine, at the time of homologous virus challenge, substantial levels of antibodies binding to FIV core (p28) and envelope (SU and TM) antigens, as measured by ELISA (1:1000-1:50000 for anti-p28 antibodies) and immunoblot (1:50-1:5000 for anti-SU antibodies) were detected (Figs

1-3). Significant titers of neutralizing antibody (mean titers of 1:50-1:200) were also achieved, generally higher in those cats immunized with the fixed infected cell vaccine (Fig 4). A microculture modification of the virus neutralization (VN) assay using uninfected FeT1 cells (FIV susceptible feline T cells) (20) was used. Unimmunized adjuvant controls had no FIV antibody detected. Three weeks after the 2nd boost, all cats were challenged IP with 10 ID_{50} of the homologous Petaluma strain of FIV, that had previously been titrated in vivo (20). For the next 9 months the cats were monitored, as previously described, for infection and latent FIV infection, for FIV antibody response and for clinical status. The results are summarized in Table 1, Part A. All 15 cats immunized with the fixed infected cell vaccine (Group A) remained uninfected and their FIV antibody levels gradually declined over the 9 month interval following challenge. Thirteen of 15 cats immunized with the inactivated whole virus vaccine (Group B) were also completely protected against infection and showed a similar, progressive decline in FIV antibody level. The two cats (249, A40) in Group B that were not protected became persistently infected by 3 weeks after challenge and their FIV antibody levels increased and remained elevated for the next 9 months. All 10 adjuvant control cats seroconverted and became persistently infected within several weeks after challenge.

At 38 weeks after challenge the 15 uninfected cats in vaccine Group A and the 13 protected cats in vaccine Group B were boosted a third time with their respective FIV vaccines. In both groups a marked anamnestic antibody response to FIV core and envelope antigens was elicited (Figs 1-3). Anti-envelope and neutralizing antibody titers peaked higher than the levels reached after the 2nd boost (Fig 3-4). Seven weeks after the 3rd boost the 28 vaccine protected animals were challenged IP with 10 ID_{50} of the heterologous Dixon strain of FIV. This strain differs by 11% and 4%, respectively, in the outer envelope

and transmembrane envelope amino acid sequences, from the FIV Petaluma strain (Fig 5). In vivo titration of this FIV strain in 8 SPF cats (2 per dose from 10^{-1} to 10^{-4}) established that a 10^{-3} dilution of the stock represented 10 ID₅₀. Five of 5 SPF controls were readily infected with this dose of FIV Dixon strain. At 16 weeks after the heterologous FIV challenge all but one of the 28 vaccinated cats in Groups A and B remained uninfected (Table 1B) and showed a steady decline in FIV antibody titers. The single infected cat in vaccine Group A became positive for infectious FIV by 16 weeks post challenge and showed an increase in FIV antibodies. The time of virus isolation suggest that this cat was infected with FIV-Dixon instead of FIV-Petaluma, however further studies (ie. sequencing isolated virus from Cat 346) will be necessary to identify the FIV strain infecting this cat.

These results confirm and extend our previous findings (20) in showing that positive immune protection against homologous FIV challenge infection can be achieved in cats by either inactivated infected cell or cell-free virus vaccines. The increased efficacy noted in the present study (>90% vs 75% in previous study) can probably be attributed to use of a stronger immunogen, i.e., increased number of infected cells and amount of cell-free virus antigen per dose and/or stronger adjuvant preparation. The duration of protection has not yet been tested and the antibodies induced tended to decline rather rapidly, approaching baseline levels by 6-8 months after the last boost. However, the strong anamnestic neutralizing antibody response observed when the animals were re-boosted 10 months after their 2nd boost indicates that both vaccines have induced strong B cell memory. It is not known, however, if such a prompt memory cell response would protect against re-challenge infection in the absence of a recent booster immunization. It does not seem likely that this anamnestic response can be attributed to exposure of these animals to the homologous live

virus challenge given 38 weeks previously because the protection appeared sterilizing, i.e., no evidence of transient or latent infection was observed at that time or thereafter and the antibody levels following this challenge progressively declined. Not only do the current vaccines exhibit enhanced efficacy against the homologous strain of FIV (Petaluma strain) but they also protect very well against IP challenge infection with a heterologous (Dixon strain) FIV, differing by 11% in outer envelope amino acid sequences. This difference is similar to that between the San Diego (PPR), Glasgow and Petaluma strains of FIV (14,15,11). However, this difference is considerably less than the 21% and 22% differences at the external envelope sequences observed between the FIV-Petaluma and the Japanese (FIV-TM2)(10) and Maryland (FIV-MD) (12) isolates, respectively. Experiments are now underway to test for vaccine protection against FIV strains from Japan. Studies are also in progress to determine the variable and conserved regions in the FIV envelope, to map the immunodominant domains in core and envelope proteins and compare sequences among FIV isolates around the world. Preliminary results showing cross-neutralizing antibody activity between different European isolates and FIV-Petaluma (17) suggest that conserved immunodominant regions in the viral envelope may exist in the present vaccines produced from FIV-Petaluma.

The type of sterilization immunity achieved and the parameters of immune protection, including protection against heterologous virus strains, are remarkably similar between the FIV vaccines and the inactivated SIV vaccines. In both FIV and SIV macaque models the mechanism of immune protection achieved with the inactivated whole vaccines remains to be elucidated. In contrast to the HIV-1 chimpanzee model where neutralizing antibodies appear to be protective, a protective effect of neutralizing antibodies in the FIV and SIV models remains uncertain. In the SIV macaque model immunization with the

uninfected human T-cells, used to grow the virus for vaccination and challenge, conferred protection to 2 of 4 monkeys (16), and monkeys protected against SIV grown in the human cells were susceptible to SIV grown in rhesus cells (2). These results suggested that nonviral cellular antigens might be critical determinants of vaccine protection in the SIV macaque system. However, in another SIV macaque model system (SIV_{mne}) protection obtained by vaccination with recombinant SIV gp160 alone correlated with induction of neutralizing antibody in the absence of cellular antibodies (4,7). Yet other studies with recombinant SIV_{mac} env vaccine have failed to yield protective immunity, whether or not neutralizing antibodies were elicited (4). By contrast, in the FIV model the picture appears somewhat clearer in that immunization with the uninfected feline T cells used to grow the vaccine virus does not protect against challenge infection with FIV grown in the same cells (20). Nevertheless, in the FIV model the uninfected allogeneic feline T-cells can be shown to elicit a nonspecific adjuvant effect on the immune response, which does not, however, appear critical to the efficacy achieved (20). In the SIV model, protection by the inactivated whole SIV vaccines correlates more strongly with induction of anticellular antibodies than with induction of antiviral antibodies (9) whereas, in the FIV model, protection correlates at least equally well with antiviral as anticellular antibodies. The few vaccinated cats that failed to be protected against infection have lower antiviral (anti-SU and neutralizing) antibody titers prior to challenge than the protected cats but some vaccinated cats with extremely low levels of neutralizing antibodies were still protected from challenge. Positive passive immunization results using pooled sera from healthy cats infected with FIV (Petaluma strain) (neutralizing titer of 1:400) indicate that antiviral rather than anticellular antibodies are the critical determinant of immune protection (19). The role of vaccine-induced antibody dependent cell cytotoxicity and possibility of enhancing

antibodies in this system are also under investigation.

Initial efforts in Glasgow, UK, to prepare similarly efficacious inactivated whole virus vaccines, using the FIV-Glasgow strain, failed probably because little intact envelope was present in the vaccine and little or no neutralizing antibodies were induced (8). Instead of protection, enhancement of infection was elicited by the vaccine (6). Our efforts to show the presence of enhancing antibodies in the vaccine sera at different dilutions with or without complement on FeT1 cells (feline T cell line used for neutralization assay)(20) were so far unsuccessful (Hohdatsu, T., unpublished observation). Additional studies using a technique used to detect such antibodies in cats infected with feline infectious peritonitis virus (5) are in progress. Investigators from UK are now trying to confirm our positive results using the FIV Petaluma strain grown in FL-4 cells. Development of recombinant FIV envelope vaccines are also underway at both Davis, CA and Glasgow, UK.

This work was supported by NIH, grants CA-39016, AI30904, and AI27732 and research contract from Synbiotics Corporation. We thank Renee Nakamura, Violet Martinez, and Sherry Reynolds for their technical assistance.

TABLE 1

FIV VACCINE PROTECTION AGAINST HOMOLOGOUS AND HETEROLOGOUS FIV CHALLENGES

A

GROUP	-1 WK PC V ^a PCR ^b	3 WK PC V	6 WK PC V	9 WK PC V	16 WK PC V	37 WK PC V	# PROTECTED TOTAL #
GROUP A [N=15] (FIXED INFECTED-CELL VACCINE)	-	-	-	-	-	-	15/15 (100%)
GROUP B [N=15] (INACTIVATED VIRUS VACCINE)	-	249 ^c A40	249 A40	249 A40	249 A40	249 A40	13/15 (87%)
HOMOLOGOUS VIRUS CONTROL [N=10]	-	55U 55X 354	55U A42 55X JK1 354 7K2 250 DK2 9410	ALL	ALL	ALL	0/10 (0%)

B

GROUP	3 WK (48 WK) ^d V PCR	5 WK (50 WK) V PCR	8 WK (53 WK) V PCR	16 WK (61 WK) V PCR	# PROTECTED TOTAL #
GROUP A [N=15]	--	--	--	346 346	14/15 (93%)
GROUP B [N=13]	--	--	--	--	13/13 (100%)
HETEROLOGOUS VIRUS CONTROL [N=5]	2MK2 2MK2 IK2 IK3	2MK2 2MK2 IK2 IK3	2MK2 2MK2 IK2 IK3	2MK2 2MK2 IK2 IK3 IK4 2MK6	0/5 (0%)

^a Virus isolation (V); ^b Polymerase chain reaction (PCR)/Southern blot analysis; ^c Cat identification number; ^d Weeks PC with heterologous virus (Weeks PC with homologous FIV)

LEGENDS

Figure 1. Relative levels of anti-p28 antibodies in sera from vaccinated and unvaccinated-placebo cats were determined by ELISA (20) using electrophoretically-purified p28 as substrate antigen (50 μ g/well). Cats were immunized 3x with fixed infected-cell vaccine ($\blacktriangle \triangle$), inactivated whole-virus vaccine ($\blacklozenge \lozenge$), or placebo prior to homologous FIV-Petaluma challenge at 0 weeks. ELISA was performed at a serum dilution of 1:100 and the results are expressed in O.D. values (20). The dilution titer of vaccinated cats after the 2nd boost ranged from 1:1000 to 1:50000 by immunoblot analysis. Figures 1A and 1B represent results from vaccine protected ($\blacklozenge \blacktriangle$) and unprotected ($\lozenge \triangle$) cats, respectively, compared to the results from placebo cats infected with homologous (\blacksquare) and heterologous (\square) FIV. Cats A40 and 249 (Fig. 1B, \lozenge) became virus isolation positive at 3 weeks pc with homologous virus. All vaccine-protected cats (Fig. 1A, $\blacklozenge \blacktriangle$) and 5 naive SPF cats (Fig. 1A, 1B, \square) were boosted with their respective vaccine or placebo, respectively, at 38 weeks pc and challenged 7 weeks later with heterologous FIV-Dixon. Cat 346 (Fig. 1B, \triangle) became positive by virus isolation and PCR at 16 weeks after heterologous FIV challenge (or 61 weeks pc with homologous FIV). Five cats from each vaccinated and unvaccinated placebo groups were randomly selected to be tested for their anti-p28 antibody titers and the results are expressed as mean with standard deviation. Only those immunized cats which were virus isolation and/or PCR positive were specifically tested for the antibody titers.

Figure 2. Vaccinated and unvaccinated placebo cats were tested for serum antibodies to transmembrane peptide, TM(694-705), by ELISA at a serum dilution of 1:500. The amino acid sequence of this TM peptide is QELGCNQNQFFC. All vaccine protected

cats are shown in Figure 2A along with the two placebo groups infected with either homologous FIV-Petaluma (■) or heterologous FIV-Dixon (□). Cats A40 and 249 (Fig. 2A, ◇) from the inactivated whole-virus group and Cat 346 (Fig. 2B, △) from the fixed infected-cell group were unprotected from FIV challenge.

Figure 3. Vaccinated and unvaccinated placebo cats were tested for serum antibodies to viral surface envelope peptide, SU(595-615), by ELISA at a serum dilution of 1:100. This envelope sequence also published as P100 has been shown to be immunogenic in cats infected with Petaluma isolate (1). The dilution titer of vaccinated cats after 2nd boost ranged from 1:50 to 1:5000 by immunoblot analysis. The description of the animals are the same as above two figures.

Figure 4. Virus-neutralizing antibody titers (VN titers) (20) of vaccinated and unvaccinated placebo cats were tested at -9, 0, 6, 35, 42 54, and 77 weeks after homologous FIV-Petaluma challenge ($H_oFIV \rightarrow$). Individual VN titers of six to seven cats from the fixed infected-cell (Fig. 4A) and the inactivated whole-virus (Fig. 4B) groups were randomly tested. The two unprotected cats (A40 and 249)(Fig. 4B, □--□, *--*) from homologous FIV challenge and the single unprotected cat (346)(Fig. 4A, x--x) from heterologous FIV challenge were specifically tested for VN titer. Individual VN titers of seven placebo cats challenged with homologous FIV-Petaluma (Fig. 4C, ◆◇▲△■□*) and two placebo cats challenged with heterologous FIV-Dixon (Fig. 4C, ● x) are also shown. Only vaccine-protected cats were given the 3rd vaccine boost on 38 weeks pc ($3xB \rightarrow$) and challenged at 7 weeks later with heterologous FIV-Dixon ($H_eFIV \rightarrow$). Figure D represents the mean VN titers of different groups from Figures A,B, and C and their corresponding viral status. Serum

samples were considered positive for VN titer at tested dilutions when their RT or ELISA-p28 O.D. values were less than those of the positive controls which had no serum antibodies.

Figure 5. The predicted amino acid sequence of *env* gene products from FIV-Petaluma (11), FIV-Dixon, FIV-PPR (14), and FIV-TM2 (10) on lanes 1 through 4, respectively. The six variable regions described by Phillips et al. (14) are indicated by V1 through V6. Potential glycosylation sites are underlined and cysteine residues are boxed. The putative proteolytic cleavage site (▼) separates the surface envelope protein (SU) from the transmembrane (TM) protein. The asterisk (* →) represents the location where additional glycine (G) resides for FIV-TM2.

Fig. 1

FIV-p28 ANTIBODIES




Fig. 2

FIV-TM(694-705) ANTIBODIES





Fig. 3

FIV-SU(595-615) ANTIBODIES






Fig. 4



US

[illegible]

ITGSPSCANNTCNVSVIVPDYQCYLDRVDVTWLOCKINISICLTGGKMLYNKVKTKQLSYCTDPLQIPLINVTFGPNQTCMNWTSQIQDPEIPKCGWNNQMAYYNSCKWEEAKVKFHCORTSQSPGCSWFR
 390
 Dlx
 PPR
 TM2

Pot ISSWKORNREWRPDEFESKKVKISLQCNSTKNLTFAMRSSGDYGEVTCAWIEFCGRNKSCKLHAERPRICRNVWGSNTSLIDTGCNQTKVSGANPDVCTMYSNKMYNCGLONGFTMTKVDDLLIMHFNMK

520 V3

Dlx PPR TM2

R R K E E E E] V H S YVD Q M RF T K T N A D E N I PD KNLN TNPN T W A A KA T DS IE V T T T

pet KAVEMYNIAAGNSCTSDLPSSWGVMNENINSSSSSYSGTKMACPSNRGILRNWYNPVGILRQSLQYQVVKQDPDYLVPPEVMEYKPRRKRAATHVMLAAVLSIACAGTGATAGIMVTQYHQVLATHQE

fix K

PPR

TH2 L

IM

Q

780
 V4
 1967
 1968
 1969
 1970
 1971
 1972
 1973
 1974
 1975
 1976
 1977
 1978
 1979
 1980
 1981
 1982
 1983
 1984
 1985
 1986
 1987
 1988
 1989
 1990
 1991
 1992
 1993
 1994
 1995
 1996
 1997
 1998
 1999
 2000
 2001
 2002
 2003
 2004
 2005
 2006
 2007
 2008
 2009
 2010
 2011
 2012
 2013
 2014
 2015
 2016
 2017
 2018
 2019
 2020
 2021
 2022
 2023
 2024
 2025
 2026
 2027
 2028
 2029
 2030
 2031
 2032
 2033
 2034
 2035
 2036
 2037
 2038
 2039
 2040
 2041
 2042
 2043
 2044
 2045
 2046
 2047
 2048
 2049
 2050
 2051
 2052
 2053
 2054
 2055
 2056
 2057
 2058
 2059
 2060
 2061
 2062
 2063
 2064
 2065
 2066
 2067
 2068
 2069
 2070
 2071
 2072
 2073
 2074
 2075
 2076
 2077
 2078
 2079
 2080
 2081
 2082
 2083
 2084
 2085
 2086
 2087
 2088
 2089
 2090
 2091
 2092
 2093
 2094
 2095
 2096
 2097
 2098
 2099
 2100
 2101
 2102
 2103
 2104
 2105
 2106
 2107
 2108
 2109
 2110
 2111
 2112
 2113
 2114
 2115
 2116
 2117
 2118
 2119
 2120
 2121
 2122
 2123
 2124
 2125
 2126
 2127
 2128
 2129
 2130
 2131
 2132
 2133
 2134
 2135
 2136
 2137
 2138
 2139
 2140
 2141
 2142
 2143
 2144
 2145
 2146
 2147
 2148
 2149
 2150
 2151
 2152
 2153
 2154
 2155
 2156
 2157
 2158
 2159
 2160
 2161
 2162
 2163
 2164
 2165
 2166
 2167
 2168
 2169
 2170
 2171
 2172
 2173
 2174
 2175
 2176
 2177
 2178
 2179
 2180
 2181
 2182
 2183
 2184
 2185
 2186
 2187
 2188
 2189
 2190
 2191
 2192
 2193
 2194
 2195
 2196
 2197
 2198
 2199
 2200
 2201
 2202
 2203
 2204
 2205
 2206
 2207
 2208
 2209
 2210
 2211
 2212
 2213
 2214
 2215
 2216
 2217
 2218
 2219
 2220
 2221
 2222
 2223
 2224
 2225
 2226
 2227
 2228
 2229
 2230
 2231
 2232
 2233
 2234
 2235
 2236
 2237
 2238
 2239
 2240
 2241
 2242
 2243
 2244
 2245
 2246
 2247
 2248
 2249
 2250
 2251
 2252
 2253
 2254
 2255
 2256
 2257
 2258
 2259
 2260
 2261
 2262
 2263
 2264
 2265
 2266
 2267
 2268
 2269
 2270
 2271
 2272
 2273
 2274
 2275
 2276
 2277
 2278
 2279
 2280
 2281
 2282
 2283
 2284
 2285
 2286
 2287
 2288
 2289
 2290
 2291
 2292
 2293
 2294
 2295
 2296
 2297
 2298
 2299
 2300
 2301
 2302
 2303
 2304
 2305
 2306
 2307
 2308
 2309
 2310
 2311
 2312
 2313
 2314
 2315
 2316
 2317
 2318
 2319
 2320
 2321
 2322
 2323
 2324
 2325
 2326
 2327
 2328
 2329
 2330
 2331
 2332
 2333
 2334
 2335
 2336
 2337
 2338
 2339
 2340
 2341
 2342
 2343
 2344
 2345
 2346
 2347
 2348
 2349
 2350
 2351
 2352
 2353
 2354
 2355
 2356
 2357
 2358
 2359
 2360
 2361
 2362
 2363
 2364
 2365
 2366
 2367
 2368
 2369
 2370
 2371
 2372
 2373
 2374
 2375
 2376
 2377
 2378
 2379
 2380
 2381
 2382
 2383
 2384
 2385
 2386
 2387
 2388
 2389
 2390
 2391
 2392
 2393
 2394
 2395
 2396
 2397
 2398
 2399
 2400
 2401
 2402
 2403
 2404
 2405
 2406
 2407
 2408
 2409
 2410
 2411
 2412
 2413
 2414
 2415
 2416
 2417
 2418
 2419

et
ix
PR
M2

QYKGLGGIGLGVLLILQLPTLVDCIRNCHIKKIIGYTVIAMPEVEGEETQPOMELRRNGROCSMSEKEEE

I I SV I LI S V TN V6

IDD ETV K
IDD VHLSV I

855
855
855
856

V5

